

Time-dependent correlation function of the Jordan-Wigner operator as a Fredholm determinant

M. B. Zvonarev*, V. V. Cheianov†, T. Giamarchi‡

July 31, 2009

Abstract

We calculate a correlation function of the Jordan-Wigner operator in a class of free-fermion models formulated on an infinite one-dimensional lattice. We represent this function in terms of the determinant of an integrable Fredholm operator, convenient for analytic and numerical investigations. By using Wick's theorem, we avoid the form-factor summation customarily used in literature for treating similar problems.

1 Introduction and main results

In this paper we investigate the correlation function

$$D(\lambda; n, t) = \langle T \mathcal{O}_j(t) \mathcal{O}_{j'}^\dagger(0) \rangle, \quad n = j - j' \quad (1.1)$$

of the Jordan-Wigner operator

$$\mathcal{O}_j = e^{i\lambda \mathcal{N}_j}, \quad (1.2)$$

where

$$\mathcal{N}_j = \sum_{m=-\infty}^j \varrho_m, \quad \varrho_m = c_m^\dagger c_m. \quad (1.3)$$

The operator $c_m^\dagger$ (c_m) is the fermion creation (annihilation) operator at the m -th lattice site of an infinite one-dimensional (1D) lattice. These operators obey canonical anticommutation relation, $\{c_m, c_{m'}^\dagger\} = \delta_{mm'}$. The symbol T in Eq. (1.1) stands for the time ordering. The function (1.1) is a 2π -periodic function of a real parameter λ . The time evolution of the operators in Eq. (1.1) is generated by the free-fermion Hamiltonian

$$H = \int_{-\pi}^{\pi} \frac{dk}{2\pi} \xi_k \tilde{c}_k^\dagger \tilde{c}_k, \quad \tilde{c}_k = \sum_m e^{-ikm} c_m, \quad (1.4)$$

*DPMC-MaNEP, University of Geneva, 24 quai Ernest-Ansermet, 1211 Geneva 4, Switzerland

†Department of Physics, Lancaster University, Lancaster, LA1 4YB, United Kingdom

‡DPMC-MaNEP, University of Geneva, 24 quai Ernest-Ansermet, 1211 Geneva 4, Switzerland




Figure 1: Shown is an example of the single-particle energy ξ_k as the function of quasi-momentum k in the first Brillouin zone, $-\pi \leq k \leq \pi$ (solid line). For the ground state average standing in Eq. (1.1) the k -space states between $-k_F$ and k_F are occupied with fermions. This is illustrated with the bold line.

where ξ_k is the single-particle energy as a function of the Bloch quasimomentum in the first Brillouin zone, $-\pi \leq k \leq \pi$. For simplicity we assume that the equation $\xi_k = 0$ has only two solutions $k = \pm k_F$. In this case, the ground state average $\langle \dots \rangle$ in Eq. (1.1) is taken over the Fermi sea of fermions filling the single particle states between $-k_F$ and k_F :

$$n_F(k) = \langle \tilde{c}_k^\dagger \tilde{c}_k \rangle = \begin{cases} 1, & k \in [-k_F, k_F] \\ 0, & \text{otherwise} \end{cases} . \quad (1.5)$$

An example of ξ_k satisfying Eq. (1.5) is shown in Fig. 1.

The goal of the paper is to derive a Fredholm determinant representation for the function (1.1). This representation was used in our recent paper [1] to investigate correlation functions in the 1D Bose-Hubbard model with spin. However, in Ref. [1] we neither give the explicit expressions (1.6)–(1.12) nor explain a way they can be obtained. It is important to stress that the function (1.1) appears not only in the Bose-Hubbard model with spin. In the regime of strong interparticle repulsion it is possible for a broad class of models carrying spin and charge degrees of freedom to factorize spin and charge dynamics in such a way that the charge part is fully described by the function (1.1). This was demonstrated in Refs. [2, 3, 4, 5, 6, 7, 1] by representing the original correlation function as a convolution of the charge correlation function (1.1) with some spin correlation function of the Heisenberg model (see, for example, Eq. (3.5) of Ref. [2], Eq. (8) of Ref. [3], Eq. (11) of Ref. [1], etc.). The authors [2, 3, 4, 5, 6, 7] do not give Fredholm determinant representation for the function (1.1), analyzing it by other methods. An approach complementary to the one of Refs. [1, 2, 3, 4, 5, 6, 7] is used in Refs. [8, 9]. There 1D gas of bosons (and fermions) with spin is considered in a limit of infinitely strong short-range interparticle repulsion. The explicit expressions for the correlation functions are obtained by summing up matrix elements of the transitions to the intermediate states (form-factors summation technique). The results have the form of a Fredholm determinant convolved with some weight function. When compared with the predictions of Refs. [1, 2, 3, 4, 5, 6, 7] these results can be unfolded to the form of the convolution of some spin correlation function with the function (1.1). This way one can indeed get the Fredholm determinant representation for the function (1.1). However, in our opinion, it is easier to calculate the determinant

directly, as explained in our paper. We note that correlation functions similar to the one defined by Eq. (1.1) appear in the analysis of many other important 1D systems. The number of works on the subject is large and we, unfortunately, cannot give the exhaustive list of references here. They include the studies of the Tonks-Girardeau gas [10, 11, 12, 13], impenetrable anyons [14, 15], XY Heisenberg chain [16, 17, 18, 19, 20, 21], non-linear quantum hydrodynamics [22], and the full counting statistics of quantum transport [23].

The Fredholm determinant representation for the correlation function (1.1) reads

$$D(\lambda; n, t) = \det(\hat{I} + \hat{K}), \quad (1.6)$$

where $\hat{K}$ is a linear integral operator

$$(\hat{K}f)(k) = \int_{-k_F}^{k_F} \frac{dp}{2\pi} K(k, p) f(p) \quad (1.7)$$

with the integral kernel

$$K(k, p) = 2 \frac{\ell_+(k)\ell_-(p) - \ell_-(k)\ell_+(p)}{\tan \frac{k-p}{2}} - 2i [\ell_+(k)\ell_-(p) - \ell_-(k)\ell_+(p)] \quad (1.8)$$

defined on $[-k_F, k_F] \times [-k_F, k_F]$. The functions entering this kernel are defined as follows:

$$\ell_-(k) = \frac{1}{\sqrt{2}} \frac{1}{z(k)} \quad (1.9)$$

and

$$\ell_+(k) = \frac{1}{\sqrt{2}} \left[\frac{\sin \lambda}{2} z(k) + \frac{1 - \cos \lambda}{2} \frac{E(k)}{z(k)} \right], \quad (1.10)$$

where

$$E(k) = \int_0^{2\pi} \frac{dq}{2\pi} \frac{z^2(q) - z^2(k)}{\tan \frac{q-k}{2}} \quad (1.11)$$

and

$$z(k) = e^{\frac{i}{2}(nk - \xi_k t)}. \quad (1.12)$$

The linear integral operator with the kernel (1.8) belongs to the class of integrable operators, introduced in Refs. [24, 25], see also Ref. [13] for further discussion. For the particular case of $\lambda = \pi$ the kernel (1.8) appears in Refs. [19, 20, 21] for the XY Heisenberg chain and in Ref. [12] for the impenetrable Bose gas. The integrable operator obtained in Refs. [8, 9] coincide with ours up to a finite rank operator. The continuum limit of the kernel (1.8) is encountered in the theory of impenetrable anyons [15].

In some of the above mentioned works (for example, in Ref. [16], see also Ref. [18] for the list of references and detailed discussion) Wick's theorem is used to get the determinant representation of correlation functions, in others form-factor summation technique is employed. Within the form-factor approach the calculations have to be carried out in a finite system, and the thermodynamic limit is applied to the end result only. In contrast, Wick's theorem approach makes possible to work in the thermodynamic limit from the very beginning. Though less rigorous, the latter approach enjoys the advantage of being minimalist. In particular, it avoids constructing the determinant representation for the function (1.1) in the system with a finite number of particles and, therefore, significantly simplifies the presentation as compared with the form-factor approach. This motivates the use of the Wick's theorem in the present work.




Figure 2: Shown are two sets of points: $\mathcal{S}_1$, Eq. (2.9), and $\mathcal{S}_2$, Eq. (2.10), and the enumeration of their elements used in Eq. (2.13).

2 Derivation of the results

Consider a countable set $\mathcal{M}$ of points in space-time. For brevity we shall use the notation $\varrho(x) \equiv \varrho_j(t)$, $c(x) \equiv c_j(t)$ and $c^\dagger(x) \equiv c_j^\dagger(t)$ for the operators at the point $x = (j, t)$ from $\mathcal{M}$. For an arbitrary function $\gamma : \mathcal{M} \rightarrow \mathbb{C}$ consider the average

$$\langle T \prod_{x \in \mathcal{M}} [1 + \gamma(x) \varrho(x)] \rangle = 1 + \sum_{Q=1}^{\infty} \sum_{\mathcal{M}_Q \subset \mathcal{M}} \langle T \prod_{x \in \mathcal{M}_Q} \gamma(x) c^\dagger(x) c(x) \rangle, \quad (2.1)$$

where the inner sum is taken over all distinct subsets $\mathcal{M}_Q$ of $\mathcal{M}$ containing Q elements. Applying Wick's theorem to the average on the right hand side of Eq. (2.1) we get

$$\langle T \prod_{x \in \mathcal{M}_Q} \gamma(x) c^\dagger(x) c(x) \rangle = \det M(\mathcal{M}_Q). \quad (2.2)$$

In this expression $\det M(\mathcal{M}_Q)$ denotes the determinant of a $Q \times Q$ matrix, which for a given enumeration $x_1, \dots, x_Q$ of the elements of $\mathcal{M}_Q$ is defined by

$$[M(\mathcal{M}_Q)]_{a,b} = \gamma(x_a) G(x_a - x_b), \quad a, b = 1, \dots, Q. \quad (2.3)$$

Here

$$G(x) \equiv G(j, t) = -\langle T c_j(t) c_0^\dagger \rangle, \quad G(j, 0) \equiv \lim_{\epsilon \rightarrow 0} G(j, -\epsilon) \quad (2.4)$$

is the T -ordered single-particle Green's function. Combining Eqs. (2.4) and (2.2) we find

$$\langle T \prod_{x \in \mathcal{M}} [1 + \gamma(x) \varrho(x)] \rangle = \det(\hat{I} + \hat{K}). \quad (2.5)$$

where $\hat{K}$ is a linear endomorphism of the linear space $\mathcal{V}(\mathcal{M})$ of complex-valued functions on $\mathcal{M}$:

$$(\hat{K}f)(x) = \sum_{x' \in \mathcal{M}} \gamma(x) G(x - x') f(x'), \quad f \in \mathcal{V}(\mathcal{M}). \quad (2.6)$$

Let us now apply the formula (2.5) to the correlation function (1.1). Using the identity $\varrho_j^2 = \varrho_j$ following from the anti-commutativity of fermionic fields we get

$$e^{i\lambda \varrho_j} = 1 + \gamma \varrho_j, \quad \gamma \equiv e^{i\lambda} - 1. \quad (2.7)$$

The equation (1.1) is then written as

$$D(\lambda; n, t) = \langle T \prod_{x \in \mathcal{S}_1} [1 + \gamma \varrho(x)] \prod_{x \in \mathcal{S}_2} [1 + \gamma^* \varrho(x)] \rangle, \quad (2.8)$$

where

$$\mathcal{S}_1 = \{x : x = (n + 1 - a, t), a \in \mathbb{N}^*\}, \quad (2.9)$$

$$\mathcal{S}_2 = \{x : x = (1 - a, 0), a \in \mathbb{N}^*\} \quad (2.10)$$

and

$$\mathbb{N}^* = 1, 2, 3, \dots \quad (2.11)$$

The right hand side of Eq. (2.8) can be expressed as a determinant, Eq. (2.5), where $\hat{K}$ is a linear endomorphism of $\mathcal{V}(\mathcal{S}_1 \cup \mathcal{S}_2) = \mathcal{V}(\mathcal{S}_1) \oplus \mathcal{V}(\mathcal{S}_2)$. We represent $\hat{K}$ as a 2×2 matrix

$$\hat{K} = \begin{pmatrix} \hat{K}^{11} & \hat{K}^{12} \\ \hat{K}^{21} & \hat{K}^{22} \end{pmatrix}, \quad K^{\alpha\beta} : \mathcal{V}(\mathcal{S}_\alpha) \rightarrow \mathcal{V}(\mathcal{S}_\beta), \quad \alpha, \beta = 1, 2. \quad (2.12)$$

Using the enumeration of elements of $\mathcal{S}_1$ and $\mathcal{S}_2$ defined in Eqs. (2.9) and (2.10) and shown in Fig. 2 we write a matrix element of the representation (2.12):

$$[\hat{K}]_{a,b} = \begin{pmatrix} \gamma^* G(b-a) & \gamma^* G(b-a-n, -t) \\ \gamma G(b-a+n, t) & \gamma G(b-a) \end{pmatrix}, \quad a, b \in \mathbb{N}. \quad (2.13)$$

It is convenient to extend this definition to $a, b \in \mathbb{Z}$ by introducing

$$[\hat{U}]_{a,b} = \begin{cases} K_{a+1,b+1}, & (a, b) \in \mathbb{Z}^+ \times \mathbb{Z}^+ \\ 0, & (a, b) \in (\mathbb{Z} \times \mathbb{Z}) \setminus (\mathbb{Z}^+ \times \mathbb{Z}^+) \end{cases}, \quad (2.14)$$

where

$$\mathbb{Z}^+ = 0, 1, 2, 3, \dots \quad \mathbb{Z} = 0, \pm 1, \pm 2, \pm 3, \dots \quad (2.15)$$

Since $\det(\hat{I} + \hat{U}) = \det(\hat{I} + \hat{K})$ we get for Eq. (2.8)

$$D(\lambda; n, t) = \det(\hat{I} + \hat{U}). \quad (2.16)$$

Defining the Fourier transform by

$$f(j) = \int_{-\pi}^{\pi} \frac{dk}{2\pi} e^{ikj} \tilde{f}(k), \quad \tilde{f}(k) = \sum_{j=-\infty}^{\infty} e^{-ikj} f(j) \quad (2.17)$$

and applying it to Eq. (2.14) we express $\hat{U}$ as a linear integral operator

$$(\hat{U} \tilde{f})(k) = \int_{-\pi}^{\pi} \frac{dp}{2\pi} U(k, p) \tilde{f}(p) \quad (2.18)$$

with the integral kernel

$$U(k, p) = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \sum_{a,b=0}^{\infty} e^{-iq(a-b)} e^{ika} e^{-ipb} u(q), \quad (2.19)$$

where

$$u(q) = \begin{pmatrix} \gamma^* \tilde{G}(q) & \gamma^* \tilde{G}(q, -t) e^{-inq} \\ \gamma \tilde{G}(q, t) e^{inq} & \gamma \tilde{G}(q) \end{pmatrix}. \quad (2.20)$$

The function $\tilde{G}$ entering Eq. (2.20) is the Fourier transform of Green's function (2.4):

$$\tilde{G}(q, t) = -\theta(t > 0)[1 - n_F(q)]e^{-i\xi_q t} + \theta(t < 0)n_F(q)e^{-i\xi_q t}, \quad (2.21)$$

where ξ_q is the single-particle energy, Eq. (1.4), and $n_F(q)$ is the Fermi-Dirac function (1.5). Performing the summation in Eq. (2.19) we obtain

$$U(k, p) = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \eta(k - q)\eta(q - p)u(q), \quad (2.22)$$

where

$$\eta(k) = \frac{1}{1 - e^{ik - \epsilon}} \quad (2.23)$$

and ϵ is an infinitesimal positive constant.

Next, we note that

$$u(q) = v(q) + w(q), \quad (2.24)$$

where

$$v(q) = \begin{pmatrix} 0 & 0 \\ -\gamma z^2(q) & 0 \end{pmatrix} \quad (2.25)$$

and

$$w(q) = n_F(q) \begin{pmatrix} \gamma^* & \gamma^* z^{-2}(q) \\ \gamma z^2(q) & \gamma \end{pmatrix}. \quad (2.26)$$

The function $z(q)$ is defined by Eq. (1.12). Using the decomposition (2.24) we write $\hat{U} = \hat{V} + \hat{W}$, where

$$V(k, p) = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \eta(k - q)\eta(q - p)v(q) \quad (2.27)$$

and

$$W(k, p) = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \eta(k - q)\eta(q - p)w(q). \quad (2.28)$$

It follows from Eq. (2.25) that $\hat{V}^2 = 0$. This implies

$$(\hat{I} + \hat{V})^{-1} = \hat{I} - \hat{V} \quad \text{and} \quad \det(\hat{I} + \hat{V}) = 1. \quad (2.29)$$

Using the identities (2.29) we rewrite Eq. (2.16) as

$$D(\lambda; n, t) = \det[\hat{I} + (\hat{I} - \hat{V})\hat{W}]. \quad (2.30)$$

The integral operator $\hat{W}$ can be represented as a product

$$\hat{W} = \hat{A}\hat{B}, \quad (2.31)$$

where the operators $\hat{A}$ and $\hat{B}$ possess the following integral kernels

$$A(k, q) = \eta(k - q)a(q) \quad B(q, p) = \eta(q - p)b(q) \quad (2.32)$$

and

$$a(q) = \begin{pmatrix} 0 & \gamma^* z^{-1}(q) \\ 0 & \gamma z(q) \end{pmatrix} n_F(q), \quad b(q) = \begin{pmatrix} 0 & 0 \\ z(q) & z^{-1}(q) \end{pmatrix} n_F(q). \quad (2.33)$$

Since $\det(\hat{I} + \hat{A}\hat{B}) = \det(\hat{I} + \hat{B}\hat{A})$ we rewrite equation (2.30) as

$$D(\lambda; n, t) = \det[\hat{I} + \hat{B}(\hat{I} - \hat{V})\hat{A}]. \quad (2.34)$$

It follows from Eqs. (2.32) and (2.33) that

$$\hat{B}(\hat{I} - \hat{V})\hat{A} = \begin{pmatrix} 0 & 0 \\ 0 & \hat{K} \end{pmatrix}, \quad (2.35)$$

where

$$\hat{K} = \hat{B}_{21}\hat{A}_{12} + \hat{B}_{22}\hat{A}_{22} - \hat{B}_{22}\hat{V}_{21}\hat{A}_{12}. \quad (2.36)$$

The expression on the right hand side of (2.36) is evaluated with the help of identity

$$\int_{-\pi}^{\pi} \frac{dq}{2\pi} \eta(k-q)\eta(q-p) = \eta(k-p) \quad (2.37)$$

which gives for the integral kernel of the operator $\hat{K}$ in Eq. (2.36)

$$\begin{aligned} K(k, p) &= [\gamma^* z(k) z^{-1}(p) + \gamma z^{-1}(k) z(p)] \eta(k-p) \\ &\quad + \gamma \gamma^* z^{-1}(k) z^{-1}(p) \int_{-\pi}^{\pi} \frac{dq}{2\pi} \eta(k-q)\eta(q-p) z^2(q). \end{aligned} \quad (2.38)$$

The identity

$$\eta(k-q)\eta(q-p) = \eta(k-p)[\eta(k-q) - \eta(p-q)] \quad (2.39)$$

allows us to represent the last term in (2.38) as

$$\int_{-\pi}^{\pi} \frac{dq}{2\pi} \eta(k-q)\eta(q-p) z^2(q) = \left\{ \frac{1}{2} [z^2(k) + z^2(p)] + f(k) - f(p) \right\} \eta(k-p), \quad (2.40)$$

where

$$f(k) = \frac{1}{2i} \int_{-\pi}^{\pi} \frac{dq}{2\pi} \frac{z^2(q) - z^2(k)}{\tan \frac{q-k}{2}} + \frac{1}{2} \int_{-\pi}^{\pi} \frac{dq}{2\pi} z^2(q). \quad (2.41)$$

Substituting Eq. (2.40) into (2.38) we arrive at the expressions (1.7)–(1.12) for the integral kernel of the operator (2.36). Combining this result with Eqs. (2.34) and (2.35) we arrive at Eq. (1.6).

3 Conclusion

Using Wick's theorem we derived a Fredholm determinant representation, Eqs. (1.6)–(1.12), of the time-dependent correlation function (1.1) of the Jordan-Wigner operator (1.2). Generalization to other operators of the Jordan-Wigner type, such as $c_j \mathcal{O}_j$, and $q_j \mathcal{O}_j$, is straightforward within this approach. The obtained representation can be used to investigate the charge dynamics in the antiferromagnetic [2, 3, 4, 5], spin-incoherent [8, 9, 4, 5, 26, 27, 28, 29, 30, 31, 32], and spin-polarized [1, 6, 7, 33] 1D gases. It would also be interesting to generalize the calculations given here to the finite-temperature case, in particular, in view of the current interest to the anyon problem [14, 15]. Another possible use of the determinant representation is to investigate the applicability range of the low-energy effective field theory (bosonization) description of the functions (1.1). The latter description is known [34] to suffer from Stokes-like phenomena such as the loss of

the λ -periodicity of the large n, t asymptotics of the function (1.1). A way to get a λ -periodic asymptotics could be the use of the technique based on the asymptotic solution of the matrix Riemann-Hilbert problem. This technique, developed in Refs. [35, 36, 37] proved to be very helpful (see, for example, [28, 38] and references therein) for the analysis of the asymptotic behavior of the Fredholm determinants with a kernel of the type (1.8).

Acknowledgments

The authors would like to thank V. E. Korepin and M. Pustilnik for useful comments. This work was supported in part by the Swiss National Science Foundation under MaNEP and division II and by ESF under the INSTANS program.

References

- [1] Zvonarev M B, Cheianov V V and Giamarchi T, *Spin dynamics in a one-dimensional Bose-Hubbard model*, 2008 <http://arxiv.org/abs/0811.2676>
- [2] Ogata M and Shiba H, *Bethe-ansatz wave function, momentum distribution, and spin correlation in the one-dimensional strongly correlated Hubbard model*, 1990 Phys. Rev. B, **41** 2326
- [3] Aristov D N and Grepel D R, *Quasi-one-dimensional spin fluctuations in $YBa_2Cu_3O_{6+x}$* , 1997 Phys. Rev. B **55** 11358
- [4] Matveev K A, Furusaki A and Glazman L I, *Asymmetric zero-bias anomaly for strongly interacting electrons in one dimension*, 2007 Phys. Rev. Lett. **98** 096403
- [5] Matveev K A, Furusaki A and Glazman L I, *Bosonization of strongly interacting one-dimensional electrons*, 2007 Phys. Rev. B **76** 155440
- [6] Akhanev S and Tserkovnyak Y, *Spin-charge separation in a strongly correlated spin-polarized chain*, 2007 Phys. Rev. B **76** 140408
- [7] Matveev K A and Furusaki A, *Spectral functions of strongly interacting isospin- $\frac{1}{2}$ bosons in one dimension*, 2008 Phys. Rev. Lett. **101** 170403
- [8] Izergin A G and Pronko A G, *Temperature correlators in the two-component one-dimensional gas*, 1998 Nucl. Phys. B **520** 594
- [9] Izergin A G, Pronko A G and Abarenkova N I, *Temperature correlators in the one-dimensional Hubbard model in the strong coupling limit* 1998 Phys. Lett. A **245** 537
- [10] Schultz T D, *Note on the one-dimensional gas of impenetrable point-particle bosons*, 1963 J. Math. Phys. **4** 666
- [11] Lenard A, *Momentum distribution in the ground state of the one-dimensional system of impenetrable bosons* 1964 J. Math. Phys. **5** 930
- [12] Korepin V E and Slavnov N A, *The time dependent correlation function of an impenetrable Bose gas as a Fredholm minor. I*, 1990 Commun. Math. Phys. **129** 103
- [13] Korepin V E, Bogoliubov N M and Izergin A G, *Quantum Inverse Scattering Method and Correlation Functions*, 1993 Cambridge: Cambridge University Press

- [14] Patu O I, Korepin V E and Averin D V, *One-dimensional impenetrable anyons in thermal equilibrium: I. Anyonic generalization of Lenard's formula*, 2008 J. Phys. A: Math. Theor. **41** 145006
- [15] Patu O I, Korepin V E and Averin D V, *One-dimensional impenetrable anyons in thermal equilibrium: II. Determinant representation for the dynamic correlation functions* 2008 J. Phys. A: Math. Theor. **41** 255205
- [16] Lieb E, Schultz T D and Mattis D C, *Two soluble models of an antiferromagnetic chain*, 1961 Ann. Phys. **16** 407
- [17] McCoy B M, Perk J H H and Shrock R E, *Time-dependent correlation functions of the transverse Ising chain at the critical magnetic field*, 1983 Nucl. Phys. B **220** 35
- [18] Perk J H H, Capel H W, Quispel G R W and Nijhoff F W, *Finite-temperature correlations for the Ising chain in a transverse field*, 1984 Physica A **123** 1
- [19] Colomo F, Izergin A G, Korepin V E and Tognetti V, *Correlators in the Heisenberg XX0 chain as Fredholm determinants*, 1992 Phys. Lett. A **169** 243
- [20] Colomo F, Izergin A G, Korepin V E and Tognetti V, *Temperature correlation functions in the XX0 Heisenberg chain. 1*, 1993 Theor. Math. Phys. **94** 11
- [21] Its A R, Izergin A G, Korepin V E and Slavnov N A, *Temperature Correlations of Quantum Spins*, 1993 Phys. Rev. Lett. **70** 1704
- [22] Imambekov A and Glazman L I, *Universal theory of nonlinear Luttinger liquids*, 2009 Science **323** 228
- [23] Levitov L S, Lee H and Lesovik G B, *Electron counting statistics and coherent states of electric current* 1996 J. Math. Phys. **37** 4845
- [24] Its A R, Izergin A G and Korepin V E, *Temperature correlators of the impenetrable Bose Gas as an integrable system*, 1990 Commun. Math. Phys. **129** 205
- [25] Its A R, Izergin A G, Korepin V E and Slavnov N A, *Differential equations for quantum correlation functions*, 1990 Int. J. Mod. Phys. B **4** 1003
- [26] Matveev K A, *Conductance of a quantum wire in the Wigner-crystal regime*, 2004 Phys. Rev. Lett. **92** 106801
- [27] Cheianov V V and Zvonarev M B, *Nonunitary spin-charge separation in a one-dimensional fermion gas* 2004 Phys. Rev. Lett. **92** 176401
- [28] Cheianov V V and Zvonarev M B, *Zero temperature correlation functions for the impenetrable fermion gas*, 2004 J. Phys. A: Math. Gen. **37** 2261
- [29] Matveev K A, *Conductance of a quantum wire at low electron density*, 2004 Phys. Rev. B **70** 245319
- [30] Fiete G A and Balents L, *Green's function for magnetically incoherent interacting electrons in one dimension* 2004 Phys. Rev. Lett. **93** 226401
- [31] Cheianov V V, Smith H and Zvonarev M B, *Low-temperature crossover in the momentum distribution of cold atomic gases in one dimension*, 2008 Phys. Rev. A **71** 033610
- [32] Fiete G A, *The spin-incoherent Luttinger liquid*, 2007 Rev. Mod. Phys. **79** 801
- [33] Zvonarev M B, Cheianov V V and Giamarchi T, *Spin dynamics in a one-dimensional ferromagnetic Bose gas*, 2007 Phys. Rev. Lett. **99** 240404

- [34] Aristov D N, *Bosonization for a Wigner-Jordan-like transformation: Backscattering and umklapp processes on a fictitious lattice*, 1998 Phys. Rev. B **57** 12825
- [35] Its A R, *Asymptotic behavior of the solutions to the nonlinear Schrödinger equation, and isomonodromic deformations of systems of linear differential equations*, 1981 Soviet. Math. Dokl. **24** 452
- [36] Its A R, Izergin A G and Korepin V E, *Long-distance asymptotics of temperature correlators of the impenetrable Bose gas*, 1990 Commun. Math. Phys. **130** 471
- [37] Deift P and Zhou X, *A steepest descent method for oscillatory Riemann–Hilbert problems*, 1993 Ann. of Math. **137** 295
- [38] Kitanine N, Kozłowski K K, Maillet J M, Slavnov N A and Terras V, *Riemann-Hilbert approach to a generalized sine kernel and applications*, 2008 <http://arxiv.org/abs/0805.4586>